

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

D1

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 :

A61K 31/55

A2

(11) International Publication Number:

WO 96/33723

(43) International Publication Date:

31 October 1996 (31.10.96)

(21) International Application Number: PCT/JP96/01112

(22) International Filing Date: 24 April 1996 (24.04.96)

(30) Priority Data:

9508637.7

28 April 1995 (28.04.95)

GB

(71) Applicant (for all designated States except US): FUJISAWA
PHARMACEUTICAL CO., LTD. [JP/JP]; 4-7, Doshomachi
3-chome, Chuo-ku, Osaka-shi, Osaka 541 (JP).

(72) Inventor; and


(75) Inventor/Applicant (for US only): ISHIE, Masayuki [JP/JP];
5-19-4, Shinkofudai, Toyono-cho, Toyono-gun, Osaka 563-
01 (JP).(74) Agent: SEKI, Hideo; Fujisawa Pharmaceutical Co., Ltd., Osaka
Factory, 1-6, Kashima 2-chome, Yodogawa-ku, Osaka-shi,
Osaka 532 (JP).(81) Designated States: JP, US, European patent (AT, BE, CH, DE,
DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).

Published

Without international search report and to be republished
upon receipt of that report.(54) Title: USE OF BENZAMIDE DERIVATIVES FOR TREATING AND/OR PREVENTING MENIERE'S SYNDROME AND
MOTION SICKNESS

(57) Abstract

This invention relates to use of a compound of
formula (I) wherein each symbol is as defined in the
description; or its pharmaceutically acceptable salt, for
treating and/or preventing Meniere's syndrome or motion
sickness.



BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LJ	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TC	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

- 1 -

USE OF BENZAMIDE DERIVATIVES FOR TREATING AND/OR PREVENTING
MENIERE'S SYNDROME AND MOTION SICKNESS

TECHNICAL FIELD

5 This invention relates to a new use of benzamide
derivatives which possess activities as vasopressin
antagonistic activity, vasodilating activity, hypotensive
activity, activity for inhibiting saccharide release in
liver, activity for inhibiting growth of mesangium cells,
10 water diuretic activity, platelet agglutination inhibitory
activity or oxytocin antagonistic activity, for treating
and/or preventing Meniere's syndrome (e.g. Meniere's disease,
etc.) or motion sickness.

15 BACKGROUND ART

The benzamide derivatives used in this invention are known
as described in European Patent Application Publication No. 0
620 216 that said benzamide derivatives possess activities as
vasopressin antagonistic activity, vasodilating activity,
20 hypotensive activity, activity for inhibiting growth of
mesangium cells, water diuretic activity, platelet agglutination
inhibitory activity or oxytocin antagonistic activity and are
useful in the treatment and/or prevention of hypertension, heart
failure, renal insufficiency, edema, ascites, vasopressin
25 parasecretion syndrome, hepatocirrhosis, hyponatremia,
hypokalemia, diabetic, circulation disorder, oxytocin relating
diseases [e.g. premature delivery, dysmenorrhea, endometritis,
etc.] and the like in human beings and animals.

30 DISCLOSURE OF INVENTION

This invention relates to a new use of benzamide
derivatives, which possess activities as vasopressin
antagonistic activity, vasodilating activity, hypotensive
activity, activity for inhibiting saccharide release in
35 liver, activity for inhibiting growth of mesangium cells,

- 2 -

water diuretic activity, platelet agglutination inhibitory activity or oxytocin antagonistic activity, for treating and/or preventing Meniere's syndrome (e.g. Meniere's disease, etc.) or motion sickness.


5 Accordingly, this invention provides a new use of benzamide derivatives for treating and/or preventing Meniere's syndrome (e.g. Meniere's disease, etc.) or motion sickness.

10 Further, this invention provides an agent and a pharmaceutical composition for treating and/or preventing Meniere's syndrome (e.g. Meniere's disease, etc.) or motion sickness, which comprises said benzamide derivatives.

15 Still further, this invention provides a method for treating and/or preventing Meniere's syndrome (e.g. Meniere's disease, etc.) or motion sickness, which comprises administering said benzamide derivatives to mammals.

The benzamide derivatives used in this invention can be represented by the following general formula (I).

20




25

wherein R^1 is hydrogen or lower alkyl,
 30 R^2 is hydrogen, lower alkyl, halo(lower)alkyl, halogen or lower alkoxy,
 R^3 and R^4 are each hydrogen, lower alkyl or taken together to form oxo,
 R^5 is hydrogen, halogen, nitro, hydroxy, protected
 35 hydroxy, lower alkyl or lower alkoxy optionally

- 3 -

substituted with lower alkylamino,
 R^6 is hydrogen, lower alkyl or acyl,
 A is



in which

R^7 is hydrogen; lower alkyl optionally substituted with halogen, amino, lower alkylamino, protected amino, acyl, a heterocyclic group, hydroxy or protected hydroxy; or acyl; and

R^8 and R^9 are each hydrogen, or taken together to form oxo or thioxo; or

R^7 and R^8 are taken together to form a bond; and

R^9 is lower alkylamino, N-lower alkylpiperazinyl or lower alkylthio optionally substituted with lower alkylamino;

R^{10} is hydrogen;

R^{11} is hydrogen, hydroxy, lower alkylamino or lower alkyl optionally substituted with acyl; or

R^{10} and R^{11} are taken together to form oxo or lower alkoxyimino optionally substituted with acyl;


R^{12} is lower alkyl optionally substituted with acyl; and

R^{13} is lower alkyl;

X^1 is CH or N,

X^2 is CH or N,

- 4 -



in which

R^{14} is hydrogen, halogen, hydroxy or lower alkoxy,

R^{15} is aryloxy, naphthyl, phenyl substituted with substituent(s) selected from the group consisting of lower alkyl, lower alkoxy, halogen, halo(lower)alkyl, hydroxy, amino(lower)alkyl, azido(lower)alkyl, lower alkylamino(lower)alkyl, acylamino(lower)alkyl, hydroxy(lower)alkyl, cyano and acyl, or a heterocyclic group, and

R^{16} is aryl,

and

n is 0, 1, 2 or 3,

and pharmaceutically acceptable salts thereof.

Said compound (I) and pharmaceutically acceptable salts thereof are useful in the treatment and/or prevention of Meniere's syndrome (e.g. Meniere's disease, etc.) or motion sickness in mammals.

Particulars of the various definitions mentioned in this specification and preferred examples thereof are explained in the following.

The term "lower" is intended to mean a group having 1 to 6 carbon atom(s), preferably one having 1 to 4 carbon atom(s), unless otherwise provided.

Suitable "lower alkyl" and lower alkyl moiety in the

- 5 -

terms "halo(lower)alkyl", "amino(lower)alkyl", "N-lower alkylpiperazinyl", "lower alkylthio", "N-lower alkylpiperazinylcarbonyl", "lower alkylsulfonyl", "azido(lower)alkyl", "lower alkylamino(lower)alkyl", "acylamino(lower)alkyl", "hydroxy(lower)alkyl", "lower alkylcarbamoyl", "acyl(lower)alkyl" and "lower alkylamino" may be straight or branched one such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert-butyl, pentyl, hexyl or the like, in which preferable one is C₁-C₄ alkyl such as methyl, ethyl, propyl, isobutyl or tert-butyl.

Suitable "aryl" and aryl moiety in the terms "aryloxy" and "arylsulfonyl" may be phenyl, naphthyl, phenyl substituted with lower alkyl [e.g. tolyl, xylyl, mesityl, cumenyl, di(tert-butyl)pentyl, etc.] and the like, in which preferable one is phenyl or tolyl.

Suitable "halogen" may be fluorine, chlorine, bromine and iodine, in which preferable one is fluorine or chlorine.

Suitable "lower alkoxy" and lower alkoxy moiety in the term "lower alkoxyimino" may be methoxy, ethoxy, propoxy, isopropoxy, butoxy and the like, in which preferable one is methoxy or propoxy.

Suitable "lower alkylamino" may be mono or di(lower alkyl)amino such as methylamino, ethylamino, propylamino, isopropylamino, butylamino, tert-butylamino, isobutylamino, pentylamino, hexylamino, dimethylamino, diethylamino, dipropylamino, dibutylamino, diisopropylamino, dipentylamino, dihexylamino, N-methylethylamino or the like, in which preferable one is dimethylamino.

Suitable "lower alkylamino(lower)alkyl" may be mono or di(lower alkyl)amino substituted lower alkyl such as methylaminomethyl, methylaminoethyl, methylaminopropyl, methylaminobutyl, methylaminoethyl, ethylaminomethyl, ethylaminoethyl, ethylaminopropyl, ethylaminobutyl, ethylaminoethyl, dimethylaminomethyl, dimethylaminoethyl, dimethylaminopropyl, dimethylaminobutyl, dimethylaminoethyl,

- 6 -

diethylaminomethyl, diethylaminoethyl, diethylaminopropyl, diethylaminobutyl, diethylaminohexyl or the like, in which preferable one is dimethylaminoethyl, dimethylaminopropyl or dimethylaminobutyl.

- 5 Suitable "halo(lower)alkyl" may be chloromethyl, fluoromethyl, bromomethyl, difluoromethyl, dichloromethyl, trifluoromethyl, trichloromethyl, 2-fluoroethyl and the like, in which preferable one is trifluoromethyl.

- 10 Suitable "heterocyclic group" and a heterocyclic moiety in the terms "a heterocyclic(lower)alkyl" and "a heterocycliccarbonyl" may be one containing at least one hetero atom selected from nitrogen, sulfur and oxygen atom, and may include saturated or unsaturated, monocyclic or polycyclic heterocyclic group, and preferable heterocyclic
- 15 group may be N-containing heterocyclic group such as unsaturated 3 to 6-membered heteromonocyclic group containing 1 to 4 nitrogen atoms, for example, pyrrolyl, pyrrolinyl, imidazolyl, pyrazolyl, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, triazolyl [e.g. 4H-1,2,4-triazolyl, 1H-1,2,3-
- 20 triazolyl, 2H-1,2,3-triazolyl, etc.], tetrazolyl [e.g. 1H-tetrazolyl, 2H-tetrazolyl, etc.], etc.;
- saturated 3 to 7-membered heteromonocyclic group containing 1 to 4 nitrogen atoms [e.g. pyrrolidinyl, imidazolidinyl, piperidyl, piperazinyl, homopiperazinyl, etc.];
- 25 saturated heteropolycyclic group containing 1 to 4 nitrogen atoms, for example, quinuclidinyl, etc.;
- unsaturated condensed heterocyclic group containing 1 to 5 nitrogen atoms, for example, indolyl, isoindolyl, indolizinyl, benzimidazolyl, quinolyl, isoquinolyl,
- 30 indazolyl, benzotriazolyl, tetrazolopyridazinyl [e.g. tetrazolo[1,5-b]pyridazinyl, etc.], etc.;
- unsaturated 3 to 6-membered heteromonocyclic group containing an oxygen atom, for example, pyranyl, furyl, etc.;
- unsaturated, 3 to 6-membered heteromonocyclic group
- 35 containing 1 to 2 sulfur atoms, for example, thienyl, etc.;

- 7 -

- unsaturated 3 to 6-membered heteromonocyclic group containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms, for example, oxazolyl, isoxazolyl, oxadiazolyl [e.g. 1,2,4-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,5-oxadiazolyl, etc.], etc.;
- 5 saturated 3 to 6-membered heteromonocyclic group containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms [e.g. morpholinyl, etc.];
- unsaturated condensed heterocyclic group containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms [e.g. benzofurazanyl, benzoxazolyl, benzoxadiazolyl, etc.];
- 10 unsaturated 3 to 6-membered heteromonocyclic group containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms, for example, thiazolyl, thiadiazolyl [e.g., 1,2,4-thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,5-thiadiazolyl, etc.], etc.;
- 15 saturated 3 to 6-membered heteromonocyclic group containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms [e.g., thiazolidinyl, etc.];
- unsaturated condensed heterocyclic group containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms [e.g., benzothiazolyl, benzothiadiazolyl, etc.];
- 20 unsaturated condensed heterocyclic group containing 1 to 2 oxygen atoms [e.g. benzofuranyl, benzodioxolyl, etc.] and the like.
- Said "heterocyclic group" may be substituted with lower alkyl optionally substituted with hydroxy, acyloxy, amino, protected amino, acyl, aryl or methylenedioxyphenyl; acyl or a heterocyclic group, in which preferable one is piperazinyl, N-methylpiperazinyl, N,N-dimethylpiperazinyl, N-methylhomopiperazinyl, N-(2-hydroxyethyl)piperazinyl, N-(2-acetoxyethyl)piperazinyl, N-(3-phthalimidopropyl)-piperazinyl, N-(3-aminopropyl)piperazinyl, N-(pyrrolidinylcarbonylmethyl)piperazinyl, N-(methylenedioxyphenylmethyl)piperazinyl, N-ethoxycarbonylpiperazinyl, N-carboxypiperazinyl, N-tert-butoxycarbonylpiperazinyl, N-pyridylpiperazinyl,
- 25
- 30
- 35

- 8 -

dimethylaminopiperidyl, pyrrolyl, pyridyl, piperidyl, morpholinyl or quinuclidinyl.

Suitable "acyl" and acyl moiety in the terms "acylamino(lower)alkyl" and "acyl(lower)alkyl" may be
5 carboxy, esterified carboxy, carbamoyl, lower alkylcarbamoyl, lower alkanoyl, aroyl, a heterocycliccarbonyl, lower alkylsulfonyl, arylsulfonyl and the like.

The esterified carboxy may be substituted or unsubstituted lower alkoxy carbonyl [e.g. methoxy carbonyl,
10 ethoxy carbonyl, propoxy carbonyl, butoxy carbonyl, tert-butoxy carbonyl, hexyloxy carbonyl, 2-(dimethylamino)ethoxy carbonyl, 2-iodoethoxy carbonyl, 2,2,2-trichloroethoxy carbonyl, etc.], substituted or unsubstituted aryloxy carbonyl [e.g. phenoxy carbonyl,
15 4-nitrophenoxy carbonyl, 2-naphthyloxy carbonyl, etc.], substituted or unsubstituted ar(lower)alkoxy carbonyl [e.g. benzyloxy carbonyl, phenethyloxy carbonyl, benzhydryloxy carbonyl, 4-nitrobenzyloxy carbonyl, etc.] and the like, in which preferable one is lower alkoxy carbonyl or
20 2-(dimethylamino)ethoxy carbonyl.

The lower alkylcarbamoyl may be mono or di(lower)alkylcarbamoyl such as methylcarbamoyl, ethylcarbamoyl, propylcarbamoyl, dimethylcarbamoyl, diethylcarbamoyl, N-methyl-N-ethylcarbamoyl or the like.

25 The lower alkanoyl may be substituted or unsubstituted one such as formyl, acetyl, propionyl, butyryl, isobutyryl, valeryl, isovaleryl, pivaloyl, hexanoyl, trifluoroacetyl or the like, in which preferable one is formyl or acetyl.

The aroyl may be substituted or unsubstituted one such
30 as benzoyl, naphthoyl, toluoyl, di(tert-butyl)benzoyl, tolylbenzoyl, aminobenzoyl, tolylbenzoylaminobenzoyl and the like.

The lower alkylsulfonyl may be methylsulfonyl, ethylsulfonyl, propylsulfonyl, butylsulfonyl and the like.

35 The arylsulfonyl may be substituted or unsubstituted one

- 9 -

such as phenylsulfonyl, tolylsulfonyl, dimethoxyphenylsulfonyl or the like, in which preferable one is dimethoxyphenylsulfonyl.

5 "N-Protective group" in "protected amino" may be common N-protective group such as substituted or unsubstituted lower alkanoyl [e.g. formyl, acetyl, propionyl, trifluoroacetyl, etc.], phthaloyl, lower alkoxycarbonyl [e.g. tert-butoxycarbonyl, tert-amylloxycarbonyl, etc.], substituted or unsubstituted aralkoxycarbonyl [e.g. benzyloxycarbonyl, 10 p-nitrobenzyloxycarbonyl, etc.], substituted or unsubstituted arenesulfonyl [e.g. benzenesulfonyl, tosyl, etc.], nitrophenylsulfonyl, aralkyl [e.g. trityl, benzyl, etc.] or the like, in which preferable one is phthaloyl or tert-butoxycarbonyl.

15 "Protected hydroxy" may be commonly protected hydroxy such as substituted lower alkoxy such as lower alkoxy(lower)alkoxy [e.g. methoxymethoxy, etc.], lower alkoxy(lower)alkoxy(lower)alkoxy [e.g. methoxyethoxymethoxy, etc.], substituted or unsubstituted 20 ar(lower)alkoxy [e.g. benzyloxy, nitrobenzyloxy, etc.], etc., acyloxy such as lower alkanoyloxy [e.g. acetoxy, propionyloxy, pivaloyloxy, etc.], aroyloxy [e.g. benzoyloxy, fluorene-carbonyloxy, etc.], lower alkoxycarbonyloxy [e.g. methoxycarbonyloxy, ethoxycarbonyloxy, propoxycarbonyloxy, 25 isopropoxycarbonyloxy, butoxycarbonyloxy, isobutoxycarbonyloxy, tert-butoxycarbonyloxy, pentyloxycarbonyloxy, hexyloxycarbonyloxy, etc.], substituted or unsubstituted ar(lower)alkoxycarbonyloxy [e.g. benzyloxycarbonyloxy, bromobenzyloxycarbonyloxy, etc.] etc., 30 tri(lower)alkylsilyloxy [e.g. trimethylsilyloxy, etc.] and the like.

The phenyl group for R¹⁵ may be substituted with 1 to 5 "substituent(s) as mentioned above, wherein the preferable number of the substituent(s) is 1 to 2.


- 10 -

Preferable compound (I) is one which has hydrogen for R^1 , hydrogen, lower alkyl or halogen for R^2 , hydrogen for R^3 , hydrogen for R^4 , hydrogen or lower alkoxy for R^5 ,

5 hydrogen for R^6 , $\begin{array}{c} \text{O} \\ \parallel \\ \text{-N-C-} \\ | \\ \text{R}^7 \end{array}$ (wherein R^7 is lower alkyl

optionally substituted with amino, lower alkylamino, protected amino, acyl or piperidino) or $\begin{array}{c} \text{-CH-} \\ | \\ \text{R}^{11} \end{array}$ (wherein

10 R^{11} is hydrogen, lower alkylamino or acyl(lower)alkyl) for

A, CH for X^1 , CH for X^2 ,  R^{15} (wherein R^{15} is phenyl


15 substituted with substituent(s) selected from the group consisting of lower alkyl, lower alkoxy, halogen, halo(lower)alkyl, hydroxy, amino(lower)alkyl, azido(lower)alkyl, lower alkylamino(lower)alkyl, acylamino(lower)alkyl, hydroxy(lower)alkyl, cyano and acyl) for Y, and 0, 1 or 2 for n.

More preferable compound (I) is one which has hydrogen for R^1 , hydrogen, lower alkyl or halogen for R^2 , hydrogen for R^3 , hydrogen for R^4 , hydrogen or lower alkoxy for R^5 ,

25 hydrogen for R^6 , $\begin{array}{c} \text{O} \\ \parallel \\ \text{-N-C-} \\ | \\ \text{R}^7 \end{array}$ (wherein R^7 is lower alkyl

substituted with N-lower alkylpiperazinylcarbonyl or lower alkyl substituted with di(lower)alkylamino) or $\begin{array}{c} \text{-CH-} \\ | \\ \text{R}^{11} \end{array}$

30 (wherein R^{11} is lower alkyl substituted with N-lower alkylpiperazinylcarbonyl) for A, CH for X^1 , CH for X^2 ,

 R^{15} (wherein R^{15} is phenyl substituted with lower

- 11 -

alkyl or di(lower alkyl)) for Y, and 0, 1 or 2 for n.

Most preferable compound (I) is 5-(4-[2-(4-methylphenyl)benzoylamino]benzoyl)-1-[(4-methyl-1-piperazinyl)carbonylmethyl]-1,3,4,5-tetrahydro-1,5-benzodiazepin-2(2H)-one.

Suitable pharmaceutically acceptable salts of the compound (I) are conventional non-toxic salts and include an acid addition salt such as an inorganic acid addition salt [e.g. hydrochloride, hydrobromide, sulfate, phosphate, etc.], an organic acid addition salt [e.g. formate, acetate, trifluoroacetate, maleate, tartrate, methanesulfonate, benzenesulfonate, toluenesulfonate, etc.], a metal salt such as an alkali metal salt [e.g. sodium salt, potassium salt, etc.] and an alkaline earth metal salt [e.g. calcium salt, magnesium salt, etc.] and the like.

It is to be noted that the compound (I) may include one or more stereoisomer(s) such as optical isomer(s) and geometrical isomer(s) due to asymmetric carbon atoms and double bond(s), and all of such isomers and mixture thereof are included within the scope of this invention.

Now in order to show the utility of the compound (I) and pharmaceutically acceptable salts thereof in this invention, the test data of the representative compound of this invention is shown in the following.

Effect on Endocochlear Potential

1. Test Method:

Guinea-pigs were anesthetized with sodium pentobarbital (30mg/kg, i.p.). After tracheotomy, animals were maintained under artificial respiration. The cochlea was exposed using a ventrolateral approach. Endocochlear potential (EP) was recorded by means of a glass microelectrode inserted into the scala media. The Ag/AgCl reference electrode was placed on the exposed neck muscle.

Arginine vasopressin (AVP) and the test compound were

- 12 -

dissolved in saline. AVP (10^{-5} M), the test compound (0.1mg/ml), or both of them were administered through the vertebral artery. 2 ml of each drug solution was injected over a period of 2 minutes.

5

2. Test Compound:

5-(4-[2-(4-Methylphenyl)benzoylamino]benzoyl)-
1-[(4-methyl-1-piperazinyl)carbonylmethyl]-
1,3,4,5-tetrahydro-1,5-benzodiazepine-2(2H)-one
(hereinafter referred to as Compound A)

10

3. Test Result:

Data was expressed as Change in EP (Δ EP, mV). AVP produced a large negative deflection of EP from +80 to +35mV, whereas simultaneous administration of AVP and FR161282 made no change in EP.

15

10^{-5} MAVP (Δ EP, mV)	0.1mg/ml Compound A (Δ EP, mV)	10^{-5} MAVP +0.1mg/ml Compound A (Δ EP, mV)
45	0	0

20

The result show that the compound (I) of this invention has an inhibitory activity against a negative deflection of endocochlear potential induced by arginine vasopressin and is useful for treating and/or preventing Meniere's syndrome (e.g. Meniere's disease, etc.) or motion sickness.

25

For therapeutic purpose, the compounds (I) and pharmaceutically acceptable salts thereof of the present invention can be used in a form of pharmaceutical preparation containing one of said compounds, as an active ingredient, in admixture with a pharmaceutically acceptable carrier such as an organic or inorganic solid or liquid excipient suitable for oral, parenteral or external (topical) administration.

30

35

- 13 -

The pharmaceutical preparations may be capsules, tablets, dragees, granules, solution, suspension, emulsion, ointment, or the like. If desired, there may be included in these preparations, auxiliary substances, stabilizing agents, wetting or emulsifying agents, buffers and other commonly used additives.

While the dosage of the compounds (I) will vary depending upon the age and condition of the patient, an average single dose of about 0.1 mg, 1 mg, 10 mg, 50 mg, 100 mg, 250 mg, 500 mg and 1000 mg of the compound (I) may be effective for treating and/or preventing Meniere's syndrome (e.g. Meniere's disease, etc.) or motion sickness. In general, amounts between 0.1 mg/body and about 1,000 mg/body may be administered per day.

The following example is given for the purpose of illustrating the present invention.

Example 1

(Capsule)

5-(4-[2-(4-Methylphenyl)benzoylamino]benzoyl)-
1-[(4-methyl-1-piperazinyl)carbonylmethyl]-

1,3,4,5-tetrahydro-1,5-benzodiazepine-2 (2H)-one 5 mg

Lactose 80 mg


The above-mentioned ingredients, were mixed and the mixture was encapsulated to provide the capsule.

- 14 -

CLAIMS

1. Use of a compound of the formula :

5



(I)

10

15

wherein R^1 is hydrogen or lower alkyl,

R^2 is hydrogen, lower alkyl, halo(lower)alkyl, halogen or lower alkoxy,

20


R^3 and R^4 are each hydrogen, lower alkyl or taken together to form oxo,

R^5 is hydrogen, halogen, nitro, hydroxy, protected hydroxy, lower alkyl or lower alkoxy optionally substituted with lower alkylamino,

25

R^6 is hydrogen, lower alkyl or acyl,

A is



30

in which

R^7 is hydrogen; lower alkyl optionally substituted with halogen, amino,

35

lower alkylamino, protected amino,

- 15 -

acyl, a heterocyclic group, hydroxy
or protected hydroxy; or acyl; and
 R^8 and R^9 are each hydrogen, or taken
together to form oxo or thioxo; or
 R^7 and R^8 are taken together to form a
bond; and

R^9 is lower alkylamino, N-lower
alkylpiperazinyl or lower alkylthio
optionally substituted with lower
alkylamino;

R^{10} is hydrogen;

R^{11} is hydrogen, hydroxy, lower alkylamino
or lower alkyl optionally
substituted with acyl; or


R^{10} and R^{11} are taken together to form oxo
or lower alkoxyimino optionally
substituted with acyl;

R^{12} is lower alkyl optionally substituted
with acyl; and

R^{13} is lower alkyl; .

X^1 is CH or N,

X^2 is CH or N,



in which

R^{14} is hydrogen, halogen, hydroxy or
lower alkoxy,

R^{15} is aryloxy, naphthyl, phenyl
substituted with substituent(s)
selected from the group consisting
of lower alkyl, lower alkoxy,
halogen, halo(lower)alkyl, hydroxy,

- 16 -

amino(lower)alkyl,
azido(lower)alkyl,
lower alkylamino(lower)alkyl,
acylamino(lower)alkyl,
5 hydroxy(lower)alkyl, cyano and acyl,
or a heterocyclic group, and

R¹⁶ is aryl,

and

n is 0, 1, 2 or 3,

10 or its pharmaceutically acceptable salt for treating
and/or preventing Meniere's syndrome or motion sickness.

2. A use of the compound defined in Claim 1 as an agent for
treating and/or preventing Meniere's syndrome or motion
15 sickness.

3. An agent for treating and/or preventing Meniere's
syndrome or motion sickness, which comprises the
compound defined in Claim 1.

20 4. A method for treating and/or preventing Meniere's
syndrome or motion sickness, which comprises
administering the compound defined in Claim 1 to
mammals.

25 5. A use of the compound defined in Claim 1 for
manufacturing a medicament for treating and/or
preventing Meniere's syndrome or motion sickness.

30 6. A pharmaceutical composition for treating and/or
preventing Meniere's syndrome or motion sickness, which
comprises the compound defined in Claim 1 in admixture
with a carrier or excipient.

35 7. A Process for preparing the pharmaceutical composition

- 17 -

of Claim 6, which is characterized by admixing the compound with a carrier or excipient.

5

10

15

20

25

30

35

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☒ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.